

## 1 Basic Notions

**Definition 1.** Let  $\vec{u} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$  be two vectors in  $\mathbb{R}^n$ . We define the *dot product* to be

$$\vec{u} \cdot \vec{v} = a_1b_1 + a_2b_2 + \dots + a_nb_n$$

**Example 1.** Let  $\vec{u} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$

$$\vec{u} \cdot \vec{v} = 1(4) + 2(5) + 3(6)$$

**Theorem 1.** Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^n$  and  $c \in \mathbb{R}$ .

- $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$
- $(\vec{u} + \vec{v}) \cdot \vec{w} = \vec{u} \cdot \vec{w} + \vec{v} \cdot \vec{w}$
- $(c\vec{u}) \cdot \vec{v} = c(\vec{u} \cdot \vec{v}) = \vec{u} \cdot c\vec{v}$
- $\vec{u} \cdot \vec{u} \geq 0$  and  $\vec{u} \cdot \vec{u} = 0$  if and only if  $\vec{u} = 0$ .

**Definition 2.** We define the *norm* (magnitude, length) of a vector  $\vec{u}$  to be  $\|\vec{u}\| = \sqrt{\vec{u} \cdot \vec{u}}$ . A vector with length 1 is called a *unit vector*. We define the *distance* between two vectors  $\vec{v}$  and  $\vec{u}$  is defined to be  $\|\vec{v} - \vec{u}\|$ .

Note that  $\frac{\vec{v}}{\|\vec{v}\|}$  is always a unit vector.

**Definition 3.** We say that  $\vec{u}$  is *orthogonal* (perpendicular) to  $\vec{v}$ , written  $\vec{u} \perp \vec{v}$ , if and only if  $\vec{u} \cdot \vec{v} = 0$ .

**Definition 4.** If  $W$  is a subspace, we say  $\vec{v}$  is orthogonal to  $W$ , written  $\vec{v} \perp W$  if and only if  $\vec{v} \perp \vec{w}$  for all  $\vec{w} \in W$ . We call the set of all vectors  $\vec{v} \perp W$  the *orthogonal complement* of  $W$  and denote this set  $W^\perp$ .

**Proposition 2.**  $W^\perp$  is a subspace. Moreover,  $\vec{v} \perp W$  if and only if  $\vec{v} \perp \vec{w}_i$  for  $i = 1, \dots, p$  where  $\{\vec{w}_1, \dots, \vec{w}_p\}$  is a basis for  $W$ .

**Definition 5.** A set of vectors  $\{\vec{u}_1, \dots, \vec{u}_p\}$  is orthogonal if and only if

$$\vec{u}_i \cdot \vec{u}_j = 0 \text{ when } i \neq j.$$

An *orthogonal basis* is a basis which is also an orthogonal set. An *orthonormal basis* is an orthogonal basis consisting of unit vectors.

**Theorem 3.** An orthogonal set of vectors is linearly independent. Moreover, if an orthogonal set  $\{\vec{u}_1, \dots, \vec{u}_p\}$  spans a subspace  $U$  then  $\{\vec{u}_1, \dots, \vec{u}_p\}$  is a basis for  $U$ .

Note that if  $W$  is an  $n$  dimensional vector space, any set of  $n$  orthogonal vectors is automatically a basis for  $W$ .


## 2 Orthogonal Projections

**Definition 6.** Given two vectors  $\vec{v}$  and  $\vec{u}$  we define

$$\hat{v} = \left( \frac{\vec{v} \cdot \vec{u}}{\vec{u} \cdot \vec{u}} \right) \vec{u}$$

to be the *orthogonal projection* of  $\vec{v}$  onto  $\vec{u}$ . This is sometimes denoted

$$\hat{v} = \text{proj}_{\vec{u}}(\vec{v}).$$



In the above picture the green vector  $\vec{v}$  is being projected onto the blue vector  $\vec{u}$  and the resulting red vector is the projection. Notice that the projection points in the same direction as  $\vec{u}$ .

Moreover,

$$\vec{z} = (\vec{v} - \hat{v}) \perp \vec{u}$$

and the length of  $\vec{z} = \vec{v} - \hat{v}$  is represented by the dashed line.

**Definition 7.** Let  $\vec{v}$  and  $\vec{u}$  be two vectors let  $\hat{v}$  be the projection of  $\vec{v}$  onto  $\vec{u}$ . If we define  $\vec{z} = \vec{v} - \hat{v}$  then

$$\vec{v} = \hat{v} + \vec{z}$$

where  $\hat{v}$  is parallel to  $\vec{u}$  and  $\vec{z}$  is perpendicular to  $\vec{u}$ . We call  $\hat{v}$  the component of  $\vec{v}$  parallel to  $\vec{u}$  and  $\vec{z}$  is the component of  $\vec{v}$  perpendicular to  $\vec{u}$ .

**Example 2.** Let  $\vec{v} = \begin{bmatrix} 7 \\ 6 \end{bmatrix}$  and  $\vec{u} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$ .

$$\vec{v} \cdot \vec{u} = 40, \quad \vec{u} \cdot \vec{u} = 20, \quad \hat{v} = \frac{40}{20} \vec{u} = 2\vec{u}$$

Hence we have that,

$$\vec{z} = \vec{v} - \hat{v} = \begin{bmatrix} 7 \\ 6 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

Notice that,

$$\vec{z} \cdot \vec{u} = 0$$

and

$$\vec{v} = \begin{bmatrix} 7 \\ 6 \end{bmatrix} = \begin{bmatrix} 8 \\ 4 \end{bmatrix} + \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \hat{v} + \vec{z}$$

Orthogonal basis are convenient for the following reason.

**Theorem 4.** Let  $\{\vec{u}_1, \dots, \vec{u}_p\}$  be an orthogonal basis for  $W$ . For all  $\vec{y} \in W$  we have

$$\vec{y} = c_1\vec{u}_1 + \dots + c_p\vec{u}_p \quad (1)$$

and

$$c_j = \left( \frac{\vec{y} \cdot \vec{u}_j}{\vec{u}_j \cdot \vec{u}_j} \right) \text{ for } j = 1, \dots, p \quad (2)$$


The above formulas (1) and (2) will make a bit more sense after the following definition.

**Definition 8.** Let  $W$  be a subspace of  $\mathbb{R}^n$ . Then each  $\vec{v} \in W$  can be written uniquely in the form

$$\vec{v} = \hat{v} + \vec{z}$$

where  $\hat{v} \in W$  and  $\vec{z} \in W^\perp$ . We call  $\hat{v}$  the orthogonal projection of  $\vec{v}$  onto  $W$ , sometimes denoted  $\hat{v} = \text{proj}_W(\vec{v})$ . Moreover, if  $\{\vec{u}_1, \dots, \vec{u}_p\}$  is an orthogonal basis for  $W$  then

$$\hat{v} = \left( \frac{\vec{v} \cdot \vec{u}_1}{\vec{u}_1 \cdot \vec{u}_1} \right) \vec{u}_1 + \dots + \left( \frac{\vec{v} \cdot \vec{u}_p}{\vec{u}_p \cdot \vec{u}_p} \right) \vec{u}_p$$



In the above picture,  $W$  is a 2 dimensional subspace with basis vectors  $\vec{u}_1$  and  $\vec{u}_2$  (blue vectors). The red vector,  $\hat{v}$ , is the projection of the green vector,  $\vec{v}$  onto  $W$ . Moreover, the dashed line is the length of the vector  $\vec{v} - \hat{v}$ .

**Definition 9.** The distance between a vector  $\vec{v}$  and a subspace  $W$  is given by  $\|\vec{z}\| = \|\vec{v} - \hat{v}\|$ .

Notice, (1) and (2) makes some more sense in the context of the above definition. If  $\vec{y}$  is in the subspace  $W$  then it is *equal* to its projection onto  $W$ , i.e.  $\vec{y} = \hat{y}$ .

### 3 Gram Schmidt

The Gram Schmidt process allows us to get an orthogonalized version of any basis. Essentially, at each step of the process you subtract off the projection onto a subspace. Moreover, we can rescale at each step since if  $\vec{v} \perp \vec{u}$  the  $c\vec{v} \perp \vec{u}$ .

**Theorem 5.** Given a basis  $\{\vec{x}_1, \dots, \vec{x}_p\}$  for a subspace  $W$  of  $\mathbb{R}^n$ . Define,

$$\begin{aligned}\vec{v}_1 &= \vec{x}_1 \\ \vec{v}_2 &= \vec{x}_2 - \left( \frac{\vec{x}_2 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \right) \vec{v}_1 \\ \vec{v}_3 &= \vec{x}_3 - \left( \frac{\vec{x}_3 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \right) \vec{v}_1 - \left( \frac{\vec{x}_3 \cdot \vec{v}_2}{\vec{v}_2 \cdot \vec{v}_2} \right) \vec{v}_2 \\ &\vdots \\ \vec{v}_p &= \vec{x}_p - \left( \frac{\vec{x}_p \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \right) \vec{v}_1 - \left( \frac{\vec{x}_p \cdot \vec{v}_2}{\vec{v}_2 \cdot \vec{v}_2} \right) \vec{v}_2 - \dots - \left( \frac{\vec{x}_p \cdot \vec{v}_{p-1}}{\vec{v}_{p-1} \cdot \vec{v}_{p-1}} \right) \vec{v}_{p-1}\end{aligned}$$

Then  $\{\vec{v}_1, \dots, \vec{v}_p\}$  is an orthogonal basis for  $W$  and

$$\text{span}\{\vec{v}_1, \dots, \vec{v}_k\} = \text{span}\{\vec{x}_1, \dots, \vec{x}_k\} \quad \text{for } k = 1, \dots, p.$$

**Example 3.** Let

$$\vec{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{x}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

**Step 1:**

Set  $\vec{v}_1 = \vec{x}_1$ .

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

**Step 2:**

Set  $\vec{v}_2 = \vec{x}_2 - \left( \frac{\vec{x}_2 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \right) \vec{v}_1$

$$\vec{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left( \frac{\vec{x}_2 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1} \right) \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left( \frac{3}{4} \right) \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -3/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix}.$$

Chapter 6  
Notes

**Step 3:**

Rescale

$$\vec{v}_2 = \begin{bmatrix} -3 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

**Step 4:**

Set  $\vec{v}_3 = \vec{x}_3 - \left(\frac{\vec{x}_3 \cdot \vec{v}_1}{\vec{v}_1 \cdot \vec{v}_1}\right) \vec{v}_1 - \left(\frac{\vec{x}_3 \cdot \vec{v}_2}{\vec{v}_2 \cdot \vec{v}_2}\right) \vec{v}_2$

$$\vec{v}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} - \left(\frac{2}{4}\right) \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left(\frac{2}{12}\right) \begin{bmatrix} -3 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ -2/3 \\ 1/3 \\ 1/3 \end{bmatrix}.$$

**Step 5:**

Rescale

$$\vec{v}_3 = \begin{bmatrix} 0 \\ -2 \\ 1 \\ 1 \end{bmatrix}.$$

**Result:**

We have that

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -3 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 0 \\ -2 \\ 1 \\ 1 \end{bmatrix}$$