

1. BASIC NOTIONS

Definition 1.1. A binary operation on a set S is a function $f : S \times S \rightarrow S$.

Definition 1.2. A *field* is a set \mathbb{F} together with two binary operations $+$, and \cdot called addition and multiplication (respectively) such that

1. For all $a, b, c \in \mathbb{F}$ we have

$$a + (b + c) = (a + b) + c$$

and

$$a \cdot (b \cdot c) = (a \cdot b) \cdot c.$$

2. For all $a, b \in \mathbb{F}$ we have

$$a + b = b + a$$

and

$$a \cdot b = b \cdot a.$$

3. There exists an element $0 \in \mathbb{F}$, called an additive identity, such that for all $a \in \mathbb{F}$ we have $a + 0 = a$.
4. There exists an element $1 \in \mathbb{F}$, called a multiplicative identity, such that for all $a \in \mathbb{F}$ we have $a \cdot 1 = a$.
5. For all $a \in \mathbb{F}$ there exists an element $b \in \mathbb{F}$, called an additive inverse, such that $a + b = 0$.
6. For all $a \in \mathbb{F}$ such that $a \neq 0$ there exists an element $c \in \mathbb{F}$, called a multiplicative inverse, such that $a \cdot c = 1$.
7. For all $a, b, c \in \mathbb{F}$

$$a \cdot (b + c) = (a \cdot b) + (a \cdot c)$$

Note: Fields have a unique additive and multiplicative identity denoted 0 and 1 respectively. Moreover, when the additive and multiplicative inverses exist they are unique.

Some examples: All of the following examples are with their standard operations.

1. \mathbb{Q} (rational numbers)
2. \mathbb{R} (real numbers)
3. \mathbb{C} (complex numbers)
4. $\mathbb{Z}/p\mathbb{Z}$ for p prime (Integers modulo p)

Non example: \mathbb{Z} is not a field, it lacks multiplicative inverses.

Definition 1.3. A *vector space* V over a field \mathbb{F} is a set V with two operations called *vector addition* and *scalar multiplication* where vector addition is a function $+$: $V \times V \rightarrow V$ and scalar multiplication is a function \cdot : $\mathbb{F} \times V \rightarrow V$ such that

1. For all $u, v \in V$ we have

$$u + v = v + u$$

2. For all $u, v, w \in V$ and for all $a, b \in \mathbb{F}$ we have

$$(u + v) + w = u + (v + w)$$

and

$$(ab) \cdot v = a \cdot (b \cdot v)$$

3. There exists a vector $0 \in V$, called an additive identity, such that for all $v \in V$ we have

$$v + 0 = v$$

4. For all $v \in V$ we have a vector $w \in V$, called an additive inverse, such that

$$v + w = 0$$

5. For all $v \in V$ we have

$$1 \cdot v = v$$

6. For all $a, b \in \mathbb{F}$ and for all $u, v \in V$ we have

$$a \cdot (u + v) = a \cdot u + a \cdot v$$

Some examples: All of the following examples are with their standard operations.

1. $\mathbb{F}^n = \left\{ \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} : a_i \in \mathbb{F} \right\}$ where \mathbb{F} is a field.

2. Polynomials with coefficients in a field \mathbb{F} .

3. Polynomials (with coefficients in a field \mathbb{F}) of degree $\leq n$

4. Continuous functions $f : X \rightarrow Y$, $C(X, Y)$, where X and Y are fields.

5. Functions from a field X into a field Y .

6. $\mathbb{F}^\infty = \{(a_1, a_2, a_3, \dots) : a_i \in \mathbb{F}\}$.

Proposition 1.1. *Every vector space V has a unique additive identity. The unique additive identity is denoted 0 .*

Proposition 1.2. *Every element $v \in V$ has a unique additive inverse. For all $v \in V$ its unique additive inverse is denoted $-v$.*

Proposition 1.3. *For all $v \in V$ we have $0 \cdot v = 0$.*

Proposition 1.4. *For all $a \in \mathbb{F}$ and $0 \in V$ we have $a \cdot 0 = 0$.*

Proposition 1.5. *For every $v \in V$ we have $(-1) \cdot v = -v$*

2. BASIS FOR A VECTOR SPACE

Definition 2.1. A *linear combination* of a list of vectors v_1, \dots, v_m in V is a vector of the form

$$a_1v_1 + \dots + a_mv_m$$

where $a_1, \dots, a_m \in \mathbb{F}$.

Definition 2.2. The set of all linear combinations of a list of vectors v_1, \dots, v_m in V is called the *span* of v_1, \dots, v_m denoted by $\text{span}\{v_1, \dots, v_m\}$.

$$\text{span}\{v_1, \dots, v_m\} = \{a_1v_1 + \dots + a_mv_m \mid a_i \in \mathbb{F}\}$$

Definition 2.3. If V is a vector space and $V = \text{span}\{v_1, \dots, v_m\}$ then we say that v_1, \dots, v_m span V .

Definition 2.4. We say that a vectors space is *finite dimensional* if there exists a finite list of vectors v_1, \dots, v_m such that

$$\text{span}\{v_1, \dots, v_m\} = V$$

Otherwise we say that V is *infinite dimensional*.

Definition 2.5. A list of vectors v_1, \dots, v_m in V is called *linearly independent* if the only choice of $a_1, \dots, a_m \in \mathbb{F}$ such that

$$a_1v_1 + \dots + a_mv_m = 0$$

is $a_1 = a_2 = \dots = a_m = 0$. A list is called *linearly dependent* if it is not linearly independent.

Lemma 2.6. *Suppose that v_1, \dots, v_m is a linearly dependent list in V . There exists a $j \in \{1, \dots, m\}$ such that*

- 1) $v_j \in \text{span}\{v_1, \dots, v_{j-1}\}$
- 2) $\text{span}\{v_1, \dots, v_{j-1}, v_j, v_{j+1}, \dots, v_m\} = \text{span}\{v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_m\}$

Proposition 2.1. *In a finite dimensional vector space the length of every linearly independent list of vectors is less than or equal to the length of every spanning list of vectors.*

Definition 2.7. A basis for a vector space V is a list of vectors $\{v_1, \dots, v_n\}$ such that

1. $\{v_1, \dots, v_n\}$ is linearly independent
2. $\text{span}\{v_1, \dots, v_n\} = V$.

Proposition 2.2. *A list of vectors $\{v_1, \dots, v_n\}$ in V is a basis for V if and only if every vector $v \in V$ can be written uniquely in the form*

$$v = a_1v_1 + \dots + a_nv_n$$

for some $a_1, \dots, a_n \in \mathbb{F}$.

Proposition 2.3. *Every spanning list of vectors in V can be reduced down to a basis.*

Proposition 2.4. *Every linearly independent list of vectors in V can be extended to a basis.*

Proposition 2.5. *Any two basis of a finite dimensional vector space V have the same length.*

Definition 2.8. The dimension of a finite dimensional vector space is the length of any basis of the vector space. Denoted $\dim(V)$.

Proposition 2.6. *Suppose V is finite dimensional. Every list of linearly independent vectors whose length is equal to the dimension of V is a basis.*

Proposition 2.7. *Suppose V is finite dimensional. Every spanning list vectors whose length is equal to the dimension of V is a basis.*

3. SUBSPACES

Definition 3.1. A subspace of a vector space V is a subset H such that H is a vector space under the same binary relations and field as V .

Proposition 3.1 (Subspace Test). *A subset H is a subspace of V if and only if*

1. $0 \in H$.
2. For all $u, v \in H$ we have $u + v \in H$
3. For all $u \in H$ and $a \in \mathbb{F}$ we have $au \in H$.

Proposition 3.2. *If U is a subspace of a finite dimensional vector space V then $\dim(U) \leq \dim(V)$. Moreover, $\dim(U) = \dim(V)$ if and only if $V = U$.*

Definition 3.2. Suppose U_1, \dots, U_m are subsets of V . The *sum* of U_1, \dots, U_m denoted $U_1 + \dots + U_m$ is the set of all possible sums i.e.,

$$U_1 + \dots + U_m = \{u_1 + \dots + u_m \mid u_i \in U_i, i = 1, \dots, m\}$$

Proposition 3.3. *If U_1, \dots, U_m are subspaces then so is $U_1 + \dots + U_m$.*

Definition 3.3. Suppose U_1, \dots, U_m are subspaces of V . The sum $U_1 + \dots + U_m$ is a *direct sum* if each element of $U_1 + \dots + U_m$ can be written in only one way as a sum $u_1 + \dots + u_m$ where $u_i \in U_i, i = 1, \dots, m$. The direct sum is denoted $U_1 \oplus \dots \oplus U_m$.

Proposition 3.4. *$U_1 + \dots + U_m$ is a direct sum if and only if the only way to write 0 as a sum is by taking each u_i where $i = 1, \dots, m$ to be 0.*

Proposition 3.5. *The sum of two subspaces U and W is a direct sum if and only if $U \cap W = \emptyset$.*

Proposition 3.6. *If V is a finite dimensional vector space and U is a subspace of V then there exists a W which is a subspace of V such that $V = U \oplus W$.*

3.1. Quotient Spaces.

Definition 3.4. Let V be a vectors space and U a subspace. For every $v \in V$ define

$$v + U = \{v + u \mid u \in U\}$$

and

$$V/U = \{v + U \mid v \in V\}$$

Proposition 3.7. *Let V be a vectorspace, U a subspace and $v, w \in V$. The following are equivalent.*

- (a) $v - w \in U$
- (b) $v + U = w + U$
- (c) $(v + U) \cap (w + U) \neq \emptyset$

Proposition 3.8. *Let V be a vectorspace, U a subspace, $\lambda \in \mathbb{F}$, and $v, w \in V$ The set V/U is a vector space with the following operations:*

$$(v + U) + (w + U) = (v + w) + U$$

$$\lambda(v + U) = (\lambda v) + U$$

Proposition 3.9. *Let V be a finite dimensional vector space and U be a subspace.*

$$\dim(V/U) = \dim(V) - \dim(U)$$

4. LINEAR MAPS

Definition 4.1. A linear map from $V(\mathbb{F})$ to $W(\mathbb{F})$ is a function $T : V \rightarrow W$ such that

$$T(\lambda x + y) = \lambda T(x) + T(y)$$

for every $x, y \in V$ and $\lambda \in \mathbb{F}$. Denote the set of all linear maps from V to W as $\mathcal{L}(V, W)$.

Proposition 4.1. Suppose $\{v_1, \dots, v_n\}$ is a basis of V and $w_1, \dots, w_n \in W$. There is a unique linear map $T : V \rightarrow W$ such that

$$T(v_j) = w_j \quad j = 1, \dots, n.$$

Proposition 4.2. If $T : V \rightarrow W$ is linear then

$$T(0_V) = 0_W.$$

Definition 4.2. Suppose $S, T \in \mathcal{L}(V, W)$ and $\lambda \in \mathbb{F}$ define

$$(S + T)(v) = S(v) + T(v) \text{ for all } v \in V$$

and

$$(\lambda T)(v) = \lambda T(v) \text{ for all } v \in V.$$

Proposition 4.3. $\mathcal{L}(V, W)$ is a vector space with the above operations.

Definition 4.3. For $T \in \mathcal{L}(V, W)$ define the *null space* of T (or kernel of T) to be

$$\text{null}(T) = \{v \in V \mid Tv = 0\} \subseteq V$$

and define the *range* of T to be

$$\text{ran}(T) = \{Tv \mid v \in V\} \subseteq W.$$

Theorem 4.4. For $T \in \mathcal{L}(V, W)$ both $\text{null}(T)$ and $\text{ran}(T)$ are subspaces. A linear transformation is injective if and only if

$$\text{null}(T) = \{0\}.$$

A linear transformation is surjective if and only if $\text{ran}(T) = W$.

Definition 4.5. The *rank* of a linear transformation is the dimension of its range. The *nullity* of a transformation is the dimension of its null space.

Definition 4.6. A vector space isomorphism from V onto W is a bijective linear map $T : V \rightarrow W$. If there is a vector space isomorphism from V onto W we say V is isomorphic to W and write $V \cong W$.

Theorem 4.7. Let V and W be two finite dimensional vector spaces. V is isomorphic to W if and only if $\dim(V) = \dim(W)$.

Definition 4.8. A linear map $T : V \rightarrow W$ is *invertible* if there exists a map $S : W \rightarrow V$ such that

$$S \circ T = \text{Id}_V$$

$$T \circ S = \text{Id}_W$$

Theorem 4.9. A map $T : V \rightarrow W$ is invertible if and only if it is bijective.

Proposition 4.4. Suppose $T \in \mathcal{L}(V, W)$ and define

$$\tilde{T} : V/\text{null}(T) \rightarrow W$$

by

$$\tilde{T}(v + \text{null}(T)) = Tv.$$

The following hold:

- a) \tilde{T} is linear
- b) \tilde{T} is injective
- c) $\text{ran}(\tilde{T}) = \text{ran}(T)$
- d) $V/\text{null}(T) \cong \text{ran}(T)$.

Theorem 4.10 (Fundamental Theorem of Linear Maps/ Rank-Nullity). Suppose V is a finite dimensional vector space and $T \in \mathcal{L}(V, W)$. We have $\text{ran}(T)$ is finite dimensional and

$$\dim(V) = \dim(\text{ran}(T)) + \dim(\text{null}(T)).$$

Proposition 4.5. Suppose V and W are finite dimensional vector spaces. Let $T \in \mathcal{L}(V, W)$.

- a) If $\dim(V) < \dim(W)$ then T is not surjective.
- b) If $\dim(V) > \dim(W)$ then T is not injective.

Definition 4.11. A linear map $T \in \mathcal{L}(V, V)$ is called an *operator*.

Theorem 4.12. Suppose T is an operator over a vector space V . If V is finite dimension the following are equivalent:

- a) T is injective
- b) T is surjective
- c) T is bijective

4.1. The Matrix of a Linear Map and the Coordinate Transform.

Definition 4.13. Let V be an n -dimensional vector space over the field \mathbb{F} . Let $\beta = \{v_1, \dots, v_n\}$ be an *ordered* basis for V . The coordinate transform $\varphi_\beta : V \rightarrow \mathbb{F}^n$ is defined by

$$v = a_1v_1 + \dots + a_nv_n \xrightarrow{\varphi_\beta} \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$$

Denote the column vector $\varphi_\beta(v)$ by $[v]_\beta$.

Proposition 4.6. *The coordinate transform from V onto \mathbb{F}^n is a vector space isomorphism.*

Definition 4.14. Suppose $T : V \rightarrow W$ is a linear maps with $\beta = \{v_1, \dots, v_n\}$ and $\gamma = \{w_1, \dots, w_m\}$ as ordered basis for V and W respectively. Define the matrix of T , denoted by $[T]_\beta^\gamma$, by the following If

$$T(v_k) = a_{1,k}w_1 + \dots + a_{m,k}w_m$$

then the k -th column of $[T]_\beta^\gamma$ is given by $\begin{bmatrix} a_{1,k} \\ \vdots \\ a_{m,k} \end{bmatrix}$.

Theorem 4.15. *The following diagram commutes*

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \varphi_\beta \downarrow & & \downarrow \varphi_\gamma \\ \mathbb{F}^n & \xrightarrow{[T]_\beta^\gamma} & \mathbb{F}^m \end{array}$$

More specifically,

$$[T]_\beta^\gamma[v]_\beta = [T(v)]_\gamma.$$